

COMMENTARY

How citrullination invaded rheumatoid arthritis research

Walther J van Venrooij* and Ger JM Pruijn

Abstract

Citrullination and the immune response to citrullinated proteins have been fundamental for the early recognition of rheumatoid arthritis by serological tests and a better understanding of its pathophysiology. In the first years after the initial publications, the focus was on the antibodies directed to citrullinated proteins. It is now realized that citrullinating enzymes and citrullinated proteins may have important roles in the maintenance of the inflammatory processes in the joints. There is also accumulating evidence for a direct role of citrullination in tissue destruction in the rheumatoid synovium. Here we will discuss the development and importance of anti-citrullinated protein antibodies in rheumatoid arthritis as well as recent findings implicating citrullination in the pathophysiology of rheumatoid arthritis.

The first indication that patients with rheumatoid arthritis (RA) produce antibodies to a specific autoantigen was published in 1964 by two Dutch scientists, Nienhuis and Mandema. The exact nature of this antigen, the so-called perinuclear factor, remained unclear for decades. In 1978, the target of seemingly unrelated RA-specific autoantibodies (that is, keratin) was identified. Almost 15 years later, Guy Serre's group convincingly showed that both antigens were identical to the cytokeratin filament-aggregating protein filaggrin (reviewed in [1]). Our own previously published results had shown that the newly made precursor of filaggrin in cultured buccal mucosa cells (that is, profilaggrin) did not react with RA antibodies [2]. This prompted us to consider the possibility that a post-translational modification of filaggrin, absent on newly made profilaggrin, was required for the formation of the antigenic target of these antibodies. Since 1994, we

have tested several likely modifications using synthetic peptides. Indeed, citrullination, the enzymatic conversion of peptidylarginine into peptidylcitrulline, turned out to be essential to make peptides reactive with RA autoantibodies. We subsequently developed an enzyme-linked immunosorbent assay with citrullinated peptides and confirmed that the anti-peptidylcitrulline activity was specific for RA [3]. Our further work was directed to the development of the CCP2 test, using cyclic citrullinated peptides (CCPs) selected from random peptide libraries [4].

The discovery of CCP/protein as the most prominent RA-specific antigen had great impact on RA diagnostics and our understanding of RA pathophysiology. The following milestones can be noted (see [5] also).

1. After decades of intensive research by many groups, a specific diagnostic test for RA had finally been developed. The CCP2 test has a specificity of more than 95%, is very sensitive (~75%), and is still considered the gold standard in RA autoantibody testing. Since 2010, anti-citrullinated protein antibodies (ACPAs) have been included in the new American College of Rheumatology/European League Against Rheumatism classification criteria for RA.
2. Recently, an international reference preparation for ACPAs was evaluated by the International Committee for the Standardization of Autoantibodies in Rheumatic and Related Diseases [6]. It is available for the scientific community via the Centers for Disease Control and Prevention (Atlanta, GA, USA).
3. A positive CCP2 test predicts the development of RA, often years before clinical confirmation (reviewed in [5]). It appears that time to RA diagnosis is shorter in patients with high anti-CCP2 titers at enrollment as compared with those with low titers [7].

* Correspondence: wvanvenrooij@yahoo.com

Department of Biomolecular Chemistry, Institute for Molecules and Materials and Nijmegen Centre for Molecular Life Sciences, Radboud University, PO Box 9101, NL-6500 HB, Nijmegen, The Netherlands



Box 1 About Walther J van Venrooij and Ger JM Pruijn

Walther J van Venrooij received his PhD in 1968 from the University of Utrecht (The Netherlands). After that he performed a post-doctoral research study at Harvard Medical School (Boston). In 1971, he became assistant professor in biochemistry at Radboud University in Nijmegen, The Netherlands, and in 1994 he was appointed full professor in biochemistry at the same university. His research activities focused primarily on the mechanisms active in autoimmunity. In particular, he studied autoimmune diseases like rheumatoid arthritis (RA), scleroderma, systemic lupus erythematosus, and myositis. His group was the first to publish that most patients with RA have antibodies directed to citrullinated antigens (1998). Subsequently they developed the CCP2 test to measure these antibodies, which are present very early in disease. This test allows earlier diagnosis and treatment of RA and has recently been added as a new serologic criterion for classifying this disease (2010). In 2002, he received the Carol Nachmann Award for his research on autoimmune diseases. In 2004, he received the Honorary Medal from the Netherlands Society for Rheumatology.



Ger JM Pruijn received his PhD in physiological chemistry from the University of Utrecht in 1989. His interest in autoimmunity developed during his studies as a postdoctoral researcher at the University of Nijmegen and grew after obtaining a fellowship from the Royal Netherlands Academy of Arts and Sciences. In 1997, he became assistant professor in biochemistry at the University of Nijmegen and became an associate professor at the same university in 2000. Since 2006, he has been professor in biomolecular chemistry and head of the Department of Biomolecular Chemistry of the Radboud University. His research is focused on autoantibody-autoantigen systems with a special interest in the role of post-translational modifications in the autoimmune response.

based upon ACPA fine-specificity profiles is clinically meaningful.

In the last few years, it also became clear that citrullination may be instrumental to the tissue destruction processes in the synovium. ACPAs were shown to bind to osteoclast surfaces and to induce osteoclast differentiation and subsequent bone resorption [13]. This may explain why ACPA is strongly associated with bone erosion. It is tempting to speculate that the degradation of cartilage is mediated by similar mechanisms. Alternatively, this may be induced by citrullinating enzymes (peptidylarginine deiminases, or PADs), which may citrullinate proteins at the surface of cartilage, inducing their rapid degradation and destabilization of the cartilage.

Next to the antibodies, the citrullination of proteins in the synovium may lead to changes in important physiological processes. Citrullinated fibrinogen and citrullinated collagen type II are more immunogenic and arthritogenic in mouse models of arthritis, and citrullinated fibrinogen activates macrophages more than unmodified fibrinogen [10]. Recently, citrullination of fibronectin was shown to alter synovial fibroblast behavior [14]. Sokolove and colleagues [15] found that fragments of citrullinated proteins can bind to ACPAs and cause activation of macrophages and antigen-presenting cells. Ling and colleagues [16] showed that cell surface calreticulin interacts with the SE to transduce innate immune signals. Citrullination of calreticulin, an abundant process in inflamed synovial tissue, appeared to enhance its affinity for the SE, thus increasing its signaling potency significantly [16]. Interestingly, auto-citrullination of PAD4 has been reported to inactivate this enzyme and to enhance its recognition by anti-PAD4 RA sera [17]. Although not all of these data have been confirmed yet by other groups, they strongly suggest that, in addition to ACPA production, protein citrullination can affect the pathophysiology of RA at several levels.

A very interesting phenomenon that links citrullination to inflammation and autoimmunity is the formation of extracellular DNA traps, which is part of the innate immune response [18]. Several inflammatory cells externalize a meshwork of chromatin fibers, containing citrullinated histones, decorated with granule-derived (antimicrobial) proteins. Extracellular trap formation is dependent on citrullination, and proteins associated with these traps may serve as self-antigens [19]. The potential relationship between the anti-citrullinated protein response in RA and citrullination in extracellular traps remains to be investigated.

It is evident by now that the study of citrullination, ACPAs, and citrullinating enzymes becomes more and more important to understand the pathophysiology of RA. It is likely that other modifications of cellular proteins may be intrinsic factors of other autoimmune diseases. Future studies have to prove this idea.

Note: This article is part of the collection *Research through the eyes of pioneers*. Other articles in this series can be found at <http://arthritis-research.com/series/pioneers>.

Abbreviations

ACPA: Anti-citrullinated protein/peptide antibody; CCP: Cyclic citrullinated peptide; HLA: Human leukocyte antigen; Ig: Immunoglobulin; PAD: Peptidylarginine deiminase; RA: Rheumatoid arthritis; SE: Shared epitope.

Competing interests

WJV and GJM are consultants to Axis-Shield Diagnostics Ltd (Dundee, Scotland) and Euro-Diagnostica AB (Malmö, Sweden).

Published: 29 Jan 2014

References

1. Simon M, Girbal E, Sebbag M, Gomes-Daudrix V, Vincent C, Salama G, Serre G: **The cytokeratin filament-aggregating protein filaggrin is the target of the so-called "antikeratin antibodies," autoantibodies specific for rheumatoid arthritis.** *J Clin Invest* 1993, **92**:1387–1393.
2. Hoet RM, Boerbooms AM, Arends M, Ruiters DJ, van Venrooij WJ: **Antiperinuclear factor, a marker autoantibody for rheumatoid arthritis: colocalisation of the perinuclear factor and profilaggrin.** *Ann Rheum Dis* 1991, **50**:611–618.
3. Schellekens GA, de Jong BA, van den Hoogen FH, van de Putte LB, van Venrooij WJ: **Citrulline is an essential constituent of antigenic determinants recognized by rheumatoid arthritis-specific autoantibodies.** *J Clin Invest* 1998, **101**:273–281.
4. Schellekens GA, Visser H, de Jong BA, van den Hoogen FH, Hazes JM, Breedveld FC, van Venrooij WJ: **The diagnostic properties of rheumatoid arthritis antibodies recognizing a cyclic citrullinated peptide.** *Arthritis Rheum* 2000, **43**:155–163.
5. van Venrooij WJ, van Beers JJ, Pruijn GJ: **Anti-CCP antibodies: the past, the present and the future.** *Nat Rev Rheumatol* 2011, **7**:391–398.
6. Bizzaro N, Pregnotato F, van Boekel MA, Villalta D, Tozzoli R, Tonutti E, Antico A, Borghi MO, Wiik A, Meroni PL: **Preliminary evaluation of the first international reference preparation for anticitrullinated peptide antibodies.** *Ann Rheum Dis* 2012, **71**:1388–1392.
7. Bizzaro N, Bartoloni E, Morozzi G, Manganeli S, Riccieri V, Sabatini P, Filippini M, Tampoaia M, Afeltra A, Sebastiani G, Alpini C, Bini V, Bistoni O, Alunno A, Gerli R, the Forum Interdisciplinare per la Ricerca nelle Malattie Autoimmuni (FIRMA Group): **Anti-cyclic citrullinated peptide antibody titer predicts time to rheumatoid arthritis onset in patients with undifferentiated arthritis: results from a 2-year prospective study.** *Arthritis Res Ther* 2013, **15**:R16.
8. Shi J, Knevel R, Suwannaalai P, van der Linden MP, Janssen GM, van Veelen PA, Levarht NE, van der Helm-van Mil AH, Cerami A, Huizinga TW, Toes RE, Trouw LA: **Autoantibodies recognizing carbamylated proteins are present in sera of patients with rheumatoid arthritis and predict joint damage.** *Proc Natl Acad Sci U S A* 2011, **108**:17372–17377.
9. Klareskog L, Malmstrom V, Lundberg K, Padyukov L, Alfredsson L: **Smoking, citrullination and genetic variability in the immunopathogenesis of rheumatoid arthritis.** *Sem Immunol* 2011, **23**:92–98.
10. Quirke AM, Fisher BA, Kinloch AJ, Venables PJ: **Citrullination of autoantigens: upstream of TNFalpha in the pathogenesis of rheumatoid arthritis.** *FEBS Lett* 2011, **585**:3681–3688.
11. Amara K, Steen J, Murray F, Morbach H, Fernandez-Rodriguez BM, Joshua V, Engström M, Snir O, Israelsson L, Catrina AI, Wardemann H, Corti D, Meffre E, Klareskog L, Malmström V: **Monoclonal IgG antibodies generated from joint-derived B cells of RA patients have a strong bias toward citrullinated autoantigen recognition.** *J Exp Med* 2013, **210**:445–455.
12. Suwannaalai P, van de Stadt LA, Radner H, Steiner G, El-Gabalawy HS, Zijde CM, van Tol MJ, van Schaardenburg D, Huizinga TW, Toes RE, Trouw LA: **Avidity maturation of anti-citrullinated protein antibodies in rheumatoid arthritis.** *Arthritis Rheum* 2012, **64**:1323–1328.

13. Harre U, Georgess D, Bang H, Bozec A, Axmann R, Ossipova E, Jakobsson PJ, Baum W, Nimmerjahn F, Szarka E, Sarmay G, Krumbholz G, Neumann E, Toes R, Scherer HU, Catrina AJ, Klareskog L, Jurdic P, Schett G: **Induction of osteoclastogenesis and bone loss by human autoantibodies against citrullinated vimentin.** *J Clin Invest* 2012, **122**:1791–1802.
14. Shelef MA, Bennin DA, Mosher DF, Huttenlocher A: **Citrullination of fibronectin modulates synovial fibroblast behavior.** *Arthritis Res Ther* 2012, **14**:R240.
15. Sokolove J, Zhao X, Chandra PE, Robinson WH: **Immune complexes containing citrullinated fibrinogen costimulate macrophages via Toll-like receptor 4 and Fcγ receptor.** *Arthritis Rheum* 2011, **63**:53–62.
16. Ling S, Cline EN, Haug TS, Fox DA, Holoshitz J: **Citrullinated calreticulin potentiates rheumatoid arthritis shared epitope signaling.** *Arthritis Rheum* 2013, **65**:618–626.
17. Andrade F, Darrah E, Gucek M, Cole RN, Rosen A, Zhu X: **Autocitrullination of human peptidyl arginine deiminase type 4 regulates protein citrullination during cell activation.** *Arthritis Rheum* 2010, **62**:1630–1640.
18. Simon D, Simon HU, Yousefi S: **Extracellular DNA traps in allergic, infectious, and autoimmune diseases.** *Allergy* 2013, **68**:409–416.
19. Knight JS, Carmona-Rivera C, Kaplan MJ: **Proteins derived from neutrophil extracellular traps may serve as self-antigens and mediate organ damage in autoimmune diseases.** *Front Immunol* 2012, **3**:380.

10.1186/ar4458

Cite this article as: van Venrooij and Pruijn: **How citrullination invaded rheumatoid arthritis research.** *Arthritis Research & Therapy* 2014, **16**:103